arXiv:1303.0820v5 [math-ph] 29 Apr 2013

The power series expansion of Mathieu function and its integral formalism

Yoon Seok Choun[1, a)]
Baruch College
The City University of New York
Natural Science Department, A506
17 Lexington Avenue
New York, NY 10010

(Dated: 30 April 2013)

Mathieu ordinary differential equation is of Fuchsian types with the two regular and one irregular singularities. In contrast, Heun equation of Fuchsian types has the three regular and one irregular singularities. Heun equation has the four kind of confluent forms: (1) Confluent Heun (two regular and one irregular singularities), (2) Doubly confluent Heun (two irregular singularities), (3) Biconfluent Heun (one regular and one irregular singularities), (4) Triconfluent Heun equations (one regular and one irregular singularities). Mathieu equation in algebraic forms is also derived from the Confluent Heun equation by changing all coefficients $\delta = \gamma = \frac{1}{2}$, $\epsilon = 0$, $\alpha = q$ and $q = \frac{\lambda + 2q}{4}$. [see (67) in arXiv:1303.0876]

In this paper I will apply three term recurrence formula by Choun [arXiv:1303.0806] to the power series expansion in closed forms of Mathieu function (infinite series) and its integral forms including all higher terms of A_n's. One interesting observation resulting from the calculations is the fact that a Modified Bessel function recurs in each of sub-integral forms: the first sub-integral form contains zero term of $A_n's$, the second one contains one term of A_n's, the third one contains two terms of A_n's, etc. Section VI contains two additional examples of Mathieu function.

This paper is 5th out of 10 in series "Special functions and three term recurrence formula (3TRF)". See section VI for all the papers in the series. Previous paper in series deals with asymptotic behavior of Heun function and its integral formalism by Choun [arXiv:1303.0876]. The next paper in the series describes the power series expansion in closed forms of Lame function and its integral forms in the algebraic form by Choun [arXiv:1303.0873].

PACS numbers: 02.10.De, 02.30.Hq, 02.30.Ik, 02.30.Jr, 02.30.Gp, 02.30.Mv, 03.65.Ge, 03.65.-w, 03.65.Fd, 04.62.+v, 04.60.-m, 04.65.+e
Keywords: Mathieu equation, integral form, special function, three term recurrence formula, Modified Bessel function

---

[a] Electronic mail: yoon.choun@baruch.cuny.edu; ychoun@gc.cuny.edu; ychoun@gmail.com

I. INTRODUCTION

One example of three therm recursion relations is the Mathieu equation, introduced by Mathieu (1868)[14], while investigating the vibrating elliptical drumhead. Mathieu equation, known for the elliptic cylinder equation appears in diverse areas such as astronomy and physical problems involving Schrödinger equation for a periodic potentials[21], the parametric Resonance in the reheating process of universe[22], and wave equations in general relativity[18], etc. Mathieu function has been used in various areas in modern physics and mathematics.[3–11]

Unfortunately, even though Mathieu equation has been observed in various areas mentioned above, there are no power series expansion in closed forms and its integral formalism, analytically. Mathieu function has only been described in numerical approximations (Whittaker 1914[16], Frenkel and Portugal 2001[13]). Sips 1949[15], Frenkel and Portugal 2001[13] argued that it is not possible to represent analytically the Mathieu function in a simple and handy way.

I will construct the power series expansion of Mathieu function in closed forms and its integral formalism analytically using three-term recurrence formula[1] and the same method I used in analyzing Heun function.[25,26]

Mathieu equation is

$$\frac{\partial^2 y}{\partial z^2} + (\lambda - 2q\,\cos 2z)\,y = 0 \tag{1}$$

where λ and q are parameters. This is an equation with periodic-function coefficient. Mathieu equation also can be described in algebraic forms putting $x = \cos^2 z$:

$$4x(1-x)\frac{\partial^2 y}{\partial x^2} + 2(1-2x)\frac{\partial y}{\partial x} + (\lambda + 2q - 4qx)y = 0 \tag{2}$$

This equation has two regular singularities: x=0 and x=1; the other singularity x=∞ is irregular. Assume that its solution is

$$y(x) = \sum_{n=0}^{\infty} c_n x^{n+\nu} \tag{3}$$

Plug (3) into (2).

$$c_{n+1} = A_n\, c_n + B_n\, c_{n-1} \qquad ; n \geq 1 \tag{4}$$

where,

$$A_n = \frac{4(n+\nu)^2 - (\lambda + 2q)}{2(n+1+\nu)(2(n+\nu)+1)} \tag{5a}$$

$$B_n = \frac{4q}{2(n+1+\nu)(2(n+\nu)+1)} \tag{5b}$$

$$c_1 = A_0\, c_0 = \frac{1-(\lambda+2q)}{6} c_0 \tag{5c}$$

We have two indicial roots which are $\nu_1 = 0$ and $\nu_2 = \frac{1}{2}$. As we see (5b), there is no way to make B_n term terminated at certain value of n. Because the numerator of (5b) is just consist of constant q parameter. So there are only two kind of power series expansions: infinite series and the polynomial which makes A_n term terminated. In this paper I construct an analytic solution of Mathieu equation for the infinite series. And in the future I will construct its analytic solution for the polynomial which makes A_n term terminated.

II. POWER SERIES EXPANSION FOR INFINITE SERIES

In Ref. 1, the general expression of power series of $y(x)$ for infinite series is

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\
&= c_0 \Bigg\{ \sum_{i_0=0}^{\infty} \left(\prod_{i_1=0}^{i_0-1} B_{2i_1+1} \right) x^{2i_0+\lambda} + \sum_{i_0=0}^{\infty} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \sum_{i_2=i_0}^{\infty} \left(\prod_{i_3=i_0}^{i_2-1} B_{2i_3+2} \right) \right\} x^{2i_2+1+\lambda} \\
&\quad + \sum_{N=2}^{\infty} \left\{ \sum_{i_0=0}^{\infty} \left\{ A_{2i_0} \prod_{i_1=0}^{i_0-1} B_{2i_1+1} \prod_{k=1}^{N-1} \left(\sum_{i_{2k}=i_{2(k-1)}}^{\infty} A_{2i_{2k}+k} \prod_{i_{2k+1}=i_{2(k-1)}}^{i_{2k}-1} B_{2i_{2k+1}+(k+1)} \right) \right.\right. \\
&\quad \times \left.\left. \sum_{i_{2N}=i_{2(N-1)}}^{\infty} \left(\prod_{i_{2N+1}=i_{2(N-1)}}^{i_{2N}-1} B_{2i_{2N+1}+(N+1)} \right) \right\} \right\} x^{2i_{2N}+N+\lambda} \Bigg\}
\end{aligned} \tag{6}$$

Substitute (5a)-(5c) into (6). In this article Pochhammer symbol $(x)_n$ is used to represent the rising factorial: $(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)}$. And the general expression of power series of $y(x)$ for infinite series is

$$\begin{aligned} y(x) &= \sum_{n=0}^{\infty} y_n(x) = y_0(x) + y_1(x) + y_2(x) + y_3(x) + \cdots \\ &= c_0 x^{\nu} \Bigg\{ \sum_{i_0=0}^{\infty} \frac{1}{(1+\frac{\nu}{2})_{i_0}(\frac{3}{4}+\frac{\nu}{2})_{i_0}} \eta^{i_0} + x \sum_{i_0=0}^{\infty} \Bigg\{ \frac{(i_0+\frac{\nu}{2})^2 - \frac{1}{4^2}(\lambda+2q)}{(i_0+\frac{1}{2}+\frac{\nu}{2})(i_0+\frac{1}{4}+\frac{\nu}{2})} \frac{1}{(1+\frac{\nu}{2})_{i_0}(\frac{3}{4}+\frac{\nu}{2})_{i_0}} \\ &\quad \times \sum_{i_1=i_0}^{\infty} \Bigg\{ \frac{(\frac{3}{2}+\frac{\nu}{2})_{i_0}(\frac{5}{4}+\frac{\nu}{2})_{i_0}}{(\frac{3}{2}+\frac{\nu}{2})_{i_1}(\frac{5}{4}+\frac{\nu}{2})_{i_1}} \eta^{i_1} \Bigg\} \Bigg\} + \sum_{n=2}^{\infty} \Bigg\{ \sum_{i_0=0}^{\infty} \Bigg\{ \frac{(i_0+\frac{\nu}{2})^2 - \frac{1}{4^2}(\lambda+2q)}{(i_0+\frac{1}{2}+\frac{\nu}{2})(i_0+\frac{1}{4}+\frac{\nu}{2})} \frac{1}{(1+\frac{\nu}{2})_{i_0}(\frac{3}{4}+\frac{\nu}{2})_{i_0}} \\ &\quad \times \prod_{k=1}^{n-1} \Bigg\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+\frac{k}{2}+\frac{\nu}{2})^2 - \frac{1}{4^2}(\lambda+2q)}{(i_k+\frac{k}{2}+\frac{1}{2}+\frac{\nu}{2})(i_k+\frac{k}{2}+\frac{1}{4}+\frac{\nu}{2})} \frac{(1+\frac{k}{2}+\frac{\nu}{2})_{i_{k-1}}(\frac{3}{4}+\frac{k}{2}+\frac{\nu}{2})_{i_{k-1}}}{(1+\frac{k}{2}+\frac{\nu}{2})_{i_k}(\frac{3}{4}+\frac{k}{2}+\frac{\nu}{2})_{i_k}} \Bigg\} \\ &\quad \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(1+\frac{n}{2}+\frac{\nu}{2})_{i_{n-1}}(\frac{3}{4}+\frac{n}{2}+\frac{\nu}{2})_{i_{n-1}}}{(1+\frac{n}{2}+\frac{\nu}{2})_{i_n}(\frac{3}{4}+\frac{n}{2}+\frac{\nu}{2})_{i_n}} \eta^{i_n} \Bigg\} \Bigg\} x^n \Bigg\} \qquad \text{where} \quad \eta = \frac{1}{4} q x^2 \end{aligned} \tag{7}$$

As ν=0 and c_0= $\frac{1}{\Gamma(\frac{3}{4})}$ in (7),

$$\begin{aligned} y(x) &= M_f \left(x = \cos^2 z, \eta = \frac{1}{4} q x^2 \right) \\ &= \frac{1}{\Gamma\left(\frac{3}{4}\right)} \Bigg\{ \sum_{i_0=0}^{\infty} \frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}} \eta^{i_0} + x \sum_{i_0=0}^{\infty} \Bigg\{ \frac{i_0^2 - \frac{1}{4^2}(\lambda+2q)}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})} \frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}} \sum_{i_1=i_0}^{\infty} \Bigg\{ \frac{(\frac{3}{2})_{i_0}(\frac{5}{4})_{i_0}}{(\frac{3}{2})_{i_1}(\frac{5}{4})_{i_1}} \eta^{i_1} \Bigg\} \Bigg\} \\ &\quad + \sum_{n=2}^{\infty} \Bigg\{ \sum_{i_0=0}^{\infty} \Bigg\{ \frac{i_0^2 - \frac{1}{4^2}(\lambda+2q)}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})} \frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}} \\ &\quad \times \prod_{k=1}^{n-1} \Bigg\{ \sum_{i_k=i_{k-1}}^{\infty} \frac{(i_k+\frac{k}{2})^2 - \frac{1}{4^2}(\lambda+2q)}{(i_k+\frac{k}{2}+\frac{1}{2})(i_k+\frac{k}{2}+\frac{1}{4})} \frac{(1+\frac{k}{2})_{i_{k-1}}(\frac{3}{4}+\frac{k}{2})_{i_{k-1}}}{(1+\frac{k}{2})_{i_k}(\frac{3}{4}+\frac{k}{2})_{i_k}} \Bigg\} \\ &\quad \times \sum_{i_n=i_{n-1}}^{\infty} \frac{(1+\frac{n}{2})_{i_{n-1}}(\frac{3}{4}+\frac{n}{2})_{i_{n-1}}}{(1+\frac{n}{2})_{i_n}(\frac{3}{4}+\frac{n}{2})_{i_n}} \eta^{i_n} \Bigg\} \Bigg\} x^n \Bigg\} \end{aligned} \tag{8}$$

(8) is the 1^{st} kind of independent solution of Mathieu function for infinite series.

As $\nu=\frac{1}{2}$ and c_0= $\left(\frac{1}{4}q\right)^{\frac{1}{4}}/\Gamma\left(\frac{5}{4}\right)$ in (7),

$$\begin{aligned}
y(x) &= M_s\left(x=\cos^2 z, \eta=\frac{1}{4}qx^2\right)\\
&= \frac{\eta^{\frac{1}{4}}}{\Gamma\left(\frac{5}{4}\right)}\left\{\sum_{i_0=0}^{\infty}\frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}}\eta^{i_0} + x\sum_{i_0=0}^{\infty}\left\{\frac{(i_0+\frac{1}{4})^2-\frac{1}{4^2}(\lambda+2q)}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})}\frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}}\sum_{i_1=i_0}^{\infty}\left\{\frac{(\frac{7}{4})_{i_0}(\frac{3}{2})_{i_0}}{(\frac{7}{4})_{i_1}(\frac{3}{2})_{i_1}}\eta^{i_1}\right\}\right\}\right.\\
&\quad + \sum_{n=2}^{\infty}\left\{\sum_{i_0=0}^{\infty}\left\{\frac{(i_0+\frac{1}{4})^2-\frac{1}{4^2}(\lambda+2q)}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})}\frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}}\right.\right.\\
&\quad \times\prod_{k=1}^{n-1}\left\{\sum_{i_k=i_{k-1}}^{\infty}\frac{(i_k+\frac{1}{4}+\frac{k}{2})^2-\frac{1}{4^2}(\lambda+2q)}{(i_k+\frac{3}{4}+\frac{k}{2})(i_k+\frac{1}{2}+\frac{k}{2})}\frac{(\frac{5}{4}+\frac{k}{2})_{i_{k-1}}(1+\frac{k}{2})_{i_{k-1}}}{(\frac{5}{4}+\frac{k}{2})_{i_k}(1+\frac{k}{2})_{i_k}}\right\}\\
&\quad \left.\left.\left.\times\sum_{i_n=i_{n-1}}^{\infty}\frac{(\frac{5}{4}+\frac{n}{2})_{i_{n-1}}(1+\frac{n}{2})_{i_{n-1}}}{(\frac{5}{4}+\frac{n}{2})_{i_n}(1+\frac{n}{2})_{i_n}}\eta^{i_n}\right\}\right\}x^n\right\}
\end{aligned}\tag{9}$$

(9) is the 2^{nd} kind of independent solution of Mathieu function for infinite series.

III. INTEGRAL FORMALISM FOR INFINITE SERIES

There is a generalized hypergeometric function which is

$$\begin{aligned}
G_j &= \sum_{i_j=i_{j-1}}^{\infty}\frac{(1+\frac{j}{2}+\frac{\nu}{2})_{i_{j-1}}(\frac{3}{4}+\frac{j}{2}+\frac{\nu}{2})_{i_{j-1}}}{(1+\frac{j}{2}+\frac{\nu}{2})_{i_j}(\frac{3}{4}+\frac{j}{2}+\frac{\nu}{2})_{i_j}}\eta^{i_j}\\
&= \eta^{i_{j-1}}\sum_{l=0}^{\infty}\frac{1}{(i_{j-1}+\frac{3}{4}+\frac{j}{2}+\frac{\nu}{2})_l\;(i_{j-1}+1+\frac{j}{2}+\frac{\nu}{2})_l}\eta^l\\
&= \eta^{i_{j-1}}\sum_{l=0}^{\infty}\frac{B(i_{j-1}-\frac{1}{4}+\frac{j}{2}+\frac{\nu}{2},l+1)B(i_{j-1}+\frac{j}{2}+\frac{\nu}{2},l+1)}{(i_{j-1}-\frac{1}{4}+\frac{j}{2}+\frac{\nu}{2})^{-1}(i_{j-1}+\frac{j}{2}+\frac{\nu}{2})^{-1}(1)_l\;l!}\eta^l
\end{aligned}\tag{10}$$

By using integral form of beta function,

$$B\left(i_{j-1}-\frac{1}{4}+\frac{j}{2}+\frac{\nu}{2},l+1\right)=\int_0^1 dt_j\; t_j^{i_{j-1}-\frac{5}{4}+\frac{j}{2}+\frac{\nu}{2}}(1-t_j)^l \tag{11a}$$

$$B\left(i_{j-1}+\frac{j}{2}+\frac{\nu}{2},l+1\right)=\int_0^1 du_j\; u_j^{i_{j-1}-1+\frac{j}{2}+\frac{\nu}{2}}(1-u_j)^l \tag{11b}$$

Substitute (11a) and (11b) into (10). And divide $(i_{j-1}-\frac{1}{4}+\frac{j}{2}+\frac{\nu}{2})(i_{j-1}+\frac{j}{2}+\frac{\nu}{2})$ into G_j.

$$\begin{aligned}
&\frac{1}{(i_{j-1}-\frac{1}{4}+\frac{j}{2}+\frac{\nu}{2})(i_{j-1}+\frac{j}{2}+\frac{\nu}{2})}\sum_{i_j=i_{j-1}}^{\infty}\frac{(1+\frac{j}{2}+\frac{\nu}{2})_{i_{j-1}}(\frac{3}{4}+\frac{j}{2}+\frac{\nu}{2})_{i_{j-1}}}{(1+\frac{j}{2}+\frac{\nu}{2})_{i_j}(\frac{3}{4}+\frac{j}{2}+\frac{\nu}{2})_{i_j}}\eta^{i_j}\\
&= \int_0^1 dt_j\; t_j^{-\frac{5}{4}+\frac{j}{2}+\frac{\nu}{2}}\int_0^1 du_j\; u_j^{-1+\frac{j}{2}+\frac{\nu}{2}}(\eta t_j u_j)^{i_{j-1}}\sum_{l=0}^{\infty}\frac{1}{(1)_l\;l!}[\eta(1-t_j)(1-u_j)]^l
\end{aligned}\tag{12}$$

The modified Bessel function is

$$I_\alpha(x)=\sum_{l=0}^{\infty}\frac{1}{l!\;(l+\alpha)!}\left(\frac{x}{2}\right)^{2l+\alpha}=\frac{\left(\frac{x}{2}\right)^{\alpha}}{\Gamma(\frac{1}{2})\Gamma(\alpha+\frac{1}{2})}\int_{-1}^{1}dv_j\;(1-v_j^2)^{\alpha-\frac{1}{2}}e^{-xv_j} \tag{13}$$

replaced α and x by 0 and $2\sqrt{\eta(1-t_j)(1-u_j)}$ in (13).

$$I_0\left(2\sqrt{\eta(1-t_j)(1-u_j)}\right)=\sum_{l=0}^{\infty}\frac{1}{l!\;(1)_l}[\eta(1-t_j)(1-u_j)]^l=\frac{1}{\pi}\int_{-1}^{1}dv_j\;(1-v_j^2)^{-\frac{1}{2}}e^{-2\sqrt{\eta(1-t_j)(1-u_j)}\;v_j} \tag{14}$$

Substitute (14) into (12).

$$\begin{aligned}&\frac{1}{(i_{j-1}-\frac{1}{4}+\frac{j}{2}+\frac{\nu}{2})(i_{j-1}+\frac{j}{2}+\frac{\nu}{2})}\sum_{i_j=i_{j-1}}^{\infty}\frac{(1+\frac{j}{2}+\frac{\nu}{2})_{i_{j-1}}(\frac{3}{4}+\frac{j}{2}+\frac{\nu}{2})_{i_{j-1}}}{(1+\frac{j}{2}+\frac{\nu}{2})_{i_j}(\frac{3}{4}+\frac{j}{2}+\frac{\nu}{2})_{i_j}}\eta^{i_j}\\&=\int_0^1 dt_j\;t_j^{-\frac{5}{4}+\frac{j}{2}+\frac{\nu}{2}}\int_0^1 du_j\;u_j^{-1+\frac{j}{2}+\frac{\nu}{2}}I_0\left(2\sqrt{\eta(1-t_j)(1-u_j)}\right)(\eta t_j u_j)^{i_{j-1}}\end{aligned} \tag{15}$$

Defined by

$$w_{a,b}=\begin{cases}\eta\displaystyle\prod_{l=a}^{b}t_l u_l\\ \eta\;\;\text{only if } a>b\end{cases} \tag{16}$$

Substitute (15) into (7) by applying (16). The integral formalism of the function $y(x)$ for infinite series is

$$\begin{aligned}y(x)=c_0x^{\nu}\Bigg\{&\sum_{i_0=0}^{\infty}\frac{1}{(1+\frac{\nu}{2})_{i_0}(\frac{3}{4}+\frac{\nu}{2})_{i_0}}\eta^{i_0}\\&+\sum_{n=1}^{\infty}\Bigg\{\prod_{k=0}^{n-1}\Bigg\{\int_0^1 dt_{n-k}\;t_{n-k}^{-\frac{5}{4}+\frac{1}{2}(n-k+\nu)}\int_0^1 du_{n-k}\;u_{n-k}^{-1+\frac{1}{2}(n-k+\nu)}\\&\times I_0\left(2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})}\right)\\&\times\left\{w_{n-k,n}^{-\frac{1}{2}(n-1-k+\nu)}\left(w_{n-k,n}\partial_{w_{n-k,n}}\right)^2 w_{n-k,n}^{\frac{1}{2}(n-1-k+\nu)}-\frac{1}{4^2}(\lambda+2q)\right\}\\&\times\sum_{i_0=0}^{\infty}\frac{1}{(1+\frac{\nu}{2})_{i_0}(\frac{3}{4}+\frac{\nu}{2})_{i_0}}w_{1,n}^{i_0}\Bigg\}\Bigg\}x^n\Bigg\}\end{aligned} \tag{17}$$

Let $\nu = 0$ and $c_0 = \frac{1}{\Gamma(\frac{3}{4})}$ in (17).

$$\begin{aligned} y(x) &= M_f\left(x=\cos^2 z, \eta = \frac{1}{4}qx^2\right) \\ &= \frac{1}{\Gamma(\frac{3}{4})}\left\{ \sum_{i_0=0}^{\infty} \frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}}\eta^{i_0} + \sum_{n=1}^{\infty}\left\{ \prod_{k=0}^{n-1}\left\{ \int_0^1 dt_{n-k}\; t_{n-k}^{-\frac{5}{4}+\frac{1}{2}(n-k)} \int_0^1 du_{n-k}\; u_{n-k}^{-1+\frac{1}{2}(n-k)} \right.\right.\right. \\ &\quad \times I_0\left(2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})}\right) \qquad (18) \\ &\quad \left.\left.\left. \times\left\{ w_{n-k,n}^{-\frac{1}{2}(n-1-k)}\left(w_{n-k,n}\partial_{w_{n-k,n}}\right)^2 w_{n-k,n}^{\frac{1}{2}(n-1-k)} - \frac{1}{4^2}(\lambda+2q)\right\} \sum_{i_0=0}^{\infty}\frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}} w_{1,n}^{i_0}\right\}\right\} x^n \right\} \end{aligned}$$

replaced α and x by $-\frac{1}{4}$ and $2\sqrt{\eta}$ in (13).

$$I_{-\frac{1}{4}}\left(2\sqrt{\eta}\right) = \frac{\eta^{-\frac{1}{8}}}{\Gamma(\frac{3}{4})}\sum_{l=0}^{\infty}\frac{1}{l!\;(\frac{3}{4})_l}\eta^l = \frac{\eta^{-\frac{1}{8}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{4})}\int_{-1}^{1} dv_0\;(1-v_0^2)^{-\frac{3}{4}} e^{-2\sqrt{\eta}\,v_0} \qquad (19)$$

Similarly,

$$I_{-\frac{1}{4}}\left(2\sqrt{w_{1,n}}\right) = \frac{w_{1,n}^{-\frac{1}{8}}}{\Gamma(\frac{3}{4})}\sum_{l=0}^{\infty}\frac{1}{l!\;(\frac{3}{4})_l}w_{1,n}^l = \frac{w_{1,n}^{-\frac{1}{8}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{4})}\int_{-1}^{1} dv_0\;(1-v_0^2)^{-\frac{3}{4}} e^{-2\sqrt{w_{1,n}}\,v_0} \qquad (20)$$

Substitute (19) and (20) into (18).

$$\begin{aligned} y(x) &= M_f\left(x=\cos^2 z, \eta = \frac{1}{4}qx^2\right) \\ &= \eta^{\frac{1}{8}} I_{-\frac{1}{4}}\left(2\sqrt{\eta}\right) + \sum_{n=1}^{\infty}\left\{ \prod_{k=0}^{n-1}\left\{ \int_0^1 dt_{n-k}\; t_{n-k}^{-\frac{5}{4}+\frac{1}{2}(n-k)} \int_0^1 du_{n-k}\; u_{n-k}^{-1+\frac{1}{2}(n-k)} \right.\right. \\ &\quad \times I_0\left(2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})}\right) \qquad (21) \\ &\quad \left.\left. \times\left\{ w_{n-k,n}^{-\frac{1}{2}(n-1-k)}\left(w_{n-k,n}\partial_{w_{n-k,n}}\right)^2 w_{n-k,n}^{\frac{1}{2}(n-1-k)} - \frac{1}{4^2}(\lambda+2q)\right\} w_{1,n}^{\frac{1}{8}} I_{-\frac{1}{4}}\left(2\sqrt{w_{1,n}}\right) \right\}\right\} x^n \end{aligned}$$

(21) is the integral formalism of the first kind of independent solution of Mathieu function for infinite series.

Let $\nu = \frac{1}{2}$ and $c_0 = \frac{1}{\Gamma(\frac{5}{4})}\left(\frac{1}{4}q\right)^{\frac{1}{4}}$ in (17).

$$\begin{aligned} y(x) &= M_s\left(x=\cos^2 z, \eta = \frac{1}{4}qx^2\right) \\ &= \frac{\eta^{\frac{1}{4}}}{\Gamma(\frac{5}{4})}\left\{ \sum_{i_0=0}^{\infty} \frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}}\eta^{i_0} + \sum_{n=1}^{\infty}\left\{ \prod_{k=0}^{n-1}\left\{ \int_0^1 dt_{n-k}\; t_{n-k}^{-1+\frac{1}{2}(n-k)} \int_0^1 du_{n-k}\; u_{n-k}^{-\frac{3}{4}+\frac{1}{2}(n-k)} \right.\right.\right. \\ &\quad \times I_0\left(2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})}\right) \qquad (22) \\ &\quad \left.\left.\left. \times\left\{ w_{n-k,n}^{-\frac{1}{2}(n-k-\frac{1}{2})}\left(w_{n-k,n}\partial_{w_{n-k,n}}\right)^2 w_{n-k,n}^{\frac{1}{2}(n-k-\frac{1}{2})} - \frac{1}{4^2}(\lambda+2q)\right\} \sum_{i_0=0}^{\infty}\frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}} w_{1,n}^{i_0}\right\}\right\} x^n \right\} \end{aligned}$$

replaced α and x by $\frac{1}{4}$ and $2\sqrt{\eta}$ in (13).

$$I_{\frac{1}{4}}\left(2\sqrt{\eta}\right) = \frac{\eta^{\frac{1}{8}}}{\Gamma(\frac{5}{4})}\sum_{l=0}^{\infty}\frac{1}{l!\;(\frac{5}{4})_l}\eta^l = \frac{\eta^{\frac{1}{8}}}{\Gamma(\frac{1}{2})\Gamma(\frac{3}{4})}\int_{-1}^{1} dv_0\;(1-v_0^2)^{-\frac{1}{4}}e^{-2\sqrt{\eta}\,v_0} \tag{23}$$

Similarly,

$$I_{\frac{1}{4}}\left(2\sqrt{w_{1,n}}\right) = \frac{w_{1,n}^{\frac{1}{8}}}{\Gamma(\frac{5}{4})}\sum_{l=0}^{\infty}\frac{1}{l!\;(\frac{5}{4})_l}w_{1,n}^l = \frac{w_{1,n}^{\frac{1}{8}}}{\Gamma(\frac{1}{2})\Gamma(\frac{3}{4})}\int_{-1}^{1} dv_0\;(1-v_0^2)^{-\frac{1}{4}}e^{-2\sqrt{w_{1,n}}\,v_0} \tag{24}$$

Substitute (23) and (24) into (22).

$$\begin{aligned}
y(x) &= M_s\left(x=\cos^2 z, \eta=\frac{1}{4}qx^2\right)\\
&= \eta^{\frac{1}{4}}\left\{\eta^{-\frac{1}{8}}I_{\frac{1}{4}}\left(2\sqrt{\eta}\right) + \sum_{n=1}^{\infty}\left\{\prod_{k=0}^{n-1}\left\{\int_0^1 dt_{n-k}\;t_{n-k}^{-1+\frac{1}{2}(n-k)}\int_0^1 du_{n-k}\;u_{n-k}^{-\frac{3}{4}+\frac{1}{2}(n-k)}\right.\right.\right.\\
&\quad\times I_0\left(2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})}\right)\\
&\quad\left.\left.\left.\times\left\{w_{n-k,n}^{-\frac{1}{2}(n-k-\frac{1}{2})}\left(w_{n-k,n}\partial_{w_{n-k,n}}\right)^2 w_{n-k,n}^{\frac{1}{2}(n-k-\frac{1}{2})} - \frac{1}{4^2}(\lambda+2q)\right\}w_{1,n}^{-\frac{1}{8}}I_{\frac{1}{4}}\left(2\sqrt{w_{1,n}}\right)\right\}\right\}x^n\right\}
\end{aligned} \tag{25}$$

(25) is the integral formalism of the second kind of independent solution of Mathieu function for infinite series. As we see (21) and (25), Integral forms of Mathieu function are consist of modified Bessel functions surprisingly. We can transform Mathieu function from these integral forms to other well-known special functions: Kummer function, Legendre function, Hypergeometric function, Laguerre function etc.

For the special case, under the condition $\lambda \gg q > 0$, the approximate solutions of the Mathieu equation in (8) and (9) are

$$\begin{aligned}
&\lim_{\lambda\gg q>0} M_f\left(x=\cos^2 z, \eta=\frac{1}{4}qx^2\right)\\
&\simeq \frac{1}{\Gamma\left(\frac{3}{4}\right)}\left\{\sum_{i_0=0}^{\infty}\left\{\frac{1}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})}\frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}}\sum_{i_1=i_0}^{\infty}\frac{(\frac{3}{2})_{i_0}(\frac{5}{4})_{i_0}}{(\frac{3}{2})_{i_1}(\frac{5}{4})_{i_1}}\eta^{i_1}\right\}\left(-\frac{\lambda}{4^2}x\right)\right.\\
&+\sum_{n=2}^{\infty}\left\{\sum_{i_0=0}^{\infty}\left\{\frac{1}{(i_0+\frac{1}{2})(i_0+\frac{1}{4})}\frac{1}{(1)_{i_0}(\frac{3}{4})_{i_0}}\right.\right.\\
&\times\prod_{k=1}^{n-1}\left(\sum_{i_k=i_{k-1}}^{\infty}\frac{1}{(i_k+\frac{k}{2}+\frac{1}{2})(i_k+\frac{k}{2}+\frac{1}{4})}\frac{(1+\frac{k}{2})_{i_{k-1}}(\frac{3}{4}+\frac{k}{2})_{i_{k-1}}}{(1+\frac{k}{2})_{i_k}(\frac{3}{4}+\frac{k}{2})_{i_k}}\right)\\
&\left.\left.\left.\times\sum_{i_n=i_{n-1}}^{\infty}\frac{(1+\frac{n}{2})_{i_{n-1}}(\frac{3}{4}+\frac{n}{2})_{i_{n-1}}}{(1+\frac{n}{2})_{i_n}(\frac{3}{4}+\frac{n}{2})_{i_n}}\eta^{i_n}\right\}\right\}\left(-\frac{\lambda}{4^2}x\right)^n\right\}
\end{aligned} \tag{26a}$$

also,

$$\begin{aligned}
&\lim_{\lambda\gg q>0} M_s\left(x=\cos^2 z, \eta=\frac{1}{4}qx^2\right)\\
&\simeq \frac{\eta^{\frac{1}{4}}}{\Gamma\left(\frac{5}{4}\right)}\Bigg\{ \sum_{i_0=0}^{\infty}\Bigg\{\frac{1}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})}\frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}}\sum_{i_1=i_0}^{\infty}\frac{(\frac{7}{4})_{i_0}(\frac{3}{2})_{i_0}}{(\frac{7}{4})_{i_1}(\frac{3}{2})_{i_1}}\eta^{i_1}\Bigg\}\left(-\frac{\lambda}{4^2}x\right)\\
&+\sum_{n=2}^{\infty}\Bigg\{\sum_{i_0=0}^{\infty}\Bigg\{\frac{1}{(i_0+\frac{3}{4})(i_0+\frac{1}{2})}\frac{1}{(\frac{5}{4})_{i_0}(1)_{i_0}}\\
&\times\prod_{k=1}^{n-1}\left(\sum_{i_k=i_{k-1}}^{\infty}\frac{1}{(i_k+\frac{3}{4}+\frac{k}{2})(i_k+\frac{1}{2}+\frac{k}{2})}\frac{(\frac{5}{4}+\frac{k}{2})_{i_{k-1}}(1+\frac{k}{2})_{i_{k-1}}}{(\frac{5}{4}+\frac{k}{2})_{i_k}(1+\frac{k}{2})_{i_k}}\right)\\
&\times\sum_{i_n=i_{n-1}}^{\infty}\frac{(\frac{5}{4}+\frac{n}{2})_{i_{n-1}}(1+\frac{n}{2})_{i_{n-1}}}{(\frac{5}{4}+\frac{n}{2})_{i_n}(1+\frac{n}{2})_{i_n}}\eta^{i_n}\Bigg\}\Bigg\}\left(-\frac{\lambda}{4^2}x\right)^n\Bigg\} \qquad (26b)
\end{aligned}$$

And integral forms of approximated Mathieu function of (26a) and (26b) are

$$\begin{aligned}
&\lim_{\lambda\gg q>0} M_f\left(x=\cos^2 z, \eta=\frac{1}{4}qx^2\right)\\
&\simeq \frac{1}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{4})}\sum_{n=1}^{\infty}\Bigg\{\prod_{k=0}^{n-1}\Bigg\{\int_0^1 dt_{n-k}\; t_{n-k}^{-\frac{5}{4}+\frac{1}{2}(n-k)}\int_0^1 du_{n-k}\; u_{n-k}^{-1+\frac{1}{2}(n-k)}\\
&\times\frac{1}{\pi}\int_{-1}^{1} dv_{n-k}\;(1-v_{n-k}^2)^{-\frac{1}{2}}e^{-2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})}\;v_{n-k}}\\
&\times\int_{-1}^{1} dv_0\;(1-v_0^2)^{-\frac{3}{4}}e^{-2\sqrt{w_{1,n}}\;v_0}\Bigg\}\Bigg\}\left(-\frac{\lambda}{4^2}x\right)^n \qquad (27a)
\end{aligned}$$

and,

$$\begin{aligned}
&\lim_{\lambda\gg q>0} M_s\left(x=\cos^2 z, \eta=\frac{1}{4}qx^2\right)\\
&\simeq \frac{\eta^{\frac{1}{4}}}{\Gamma(\frac{1}{2})\Gamma(\frac{3}{4})}\sum_{n=1}^{\infty}\Bigg\{\prod_{k=0}^{n-1}\Bigg\{\int_0^1 dt_{n-k}\; t_{n-k}^{-1+\frac{1}{2}(n-k)}\int_0^1 du_{n-k}\; u_{n-k}^{-\frac{3}{4}+\frac{1}{2}(n-k)}\\
&\times\frac{1}{\pi}\int_{-1}^{1} dv_{n-k}\;(1-v_{n-k}^2)^{-\frac{1}{2}}e^{-2\sqrt{w_{n+1-k,n}(1-t_{n-k})(1-u_{n-k})}\;v_{n-k}}\\
&\times\int_{-1}^{1} dv_0\;(1-v_0^2)^{-\frac{1}{4}}e^{-2\sqrt{w_{1,n}}\;v_0}\Bigg\}\Bigg\}\left(-\frac{\lambda}{4^2}x\right)^n \qquad (27b)
\end{aligned}$$

IV. ASYMPTOTIC BEHAVIOR OF THE FUNCTION $y(x)$ AND THE BOUNDARY CONDITION FOR $x = \cos^2 z$ FOR THE CASE OF INFINITE SERIES

As $n \gg 1$, (5a) and (5b) are

$$\lim_{n \gg 1} A_n = A = 1 \tag{28a}$$

And,

$$\lim_{n \gg 1} B_n = B = \frac{q}{n^2} \tag{28b}$$

As $n \gg 1$, (28b) is extremely smaller than (28a) as it's compared to each other. Then, put (28a) with $B_n = 0$ into (4).

$$c_{n+1} \approx c_n \tag{29}$$

Plug (29) into the power series expansion where $\sum_{n=0}^{\infty} c_n x^n$, putting $c_0 = 1$ for simplicity.

$$\lim_{n \gg 1} y(x) = \sum_{n=0}^{\infty} x^n = \frac{1}{1-x} \qquad \text{where } 0 \leq x < 1 \tag{30}$$

As we see (30), $x = \cos^2 z$ can not be 1 for the case of infinite series. If it does't, the function $y(x)$ will blows up as $n \gg 1$ and $x = 1$.

I show the power series expansion in closed forms of Mathieu function for the infinite series in this paper analytically. Also, I derive integral forms of Mathieu function from power series expansion. It is quiet important that integral forms of this function, because we can investigate how this function is associated with other well known special functions; Bessel function, Kummer function, hypergeometric function, Laguerre function and etc. And we can obtain orthogonal relations, normalized physical factors and expectation values of any physical quantities from the integral forms of its function analytically.

V. APPLICATION

1. By using my methods on the above, we can apply the power series expansion of the Mathieu function and its integral forms into various modern physics areas.[3–10] For example, in general Mathieu equation arises from two-dimensional vibrational problems in elliptical coordinates with physical points of a view[17]. Its equation is derived from the Helmholtz equation in elliptic cylinder coordinates by using the method of separation of variables (see

(5)-(7) in p.610 in Ref. 17). Using the power series expansion of Mathieu equation, it might be possible to obtain specific eigenvalues for the wave equation in vibrational systems. Using the integral forms of Mathieu equation, it might be also possible to construct the normalized wave functions for the entire region.

2. In "Examples of Heun and Mathieu functions as solutions of wave equations in curved spaces"[18], "Dirac equation in the background of the Nutku helicoid metric"[19], two authors consider the Dirac equation in the background of the Nutku helicoid metric in five dimensions. They obtain the solution for the four different components by using the Newman-Penrose formalism[20](see (8a)-(8d) Ref. 19). And they separate uncoupled equations for the lower components into two ordinary differential equations: angular equation is of the Mathieu type and radial equation is of the double confluent form which can be reduced to the Mathieu equation (see (11), (13) Ref. 19). Using power series expansions and integral forms of Mathieu equation, it might be possible to obtain eigenvalues and normalized wave functions at various regions. In Ref. 28, I will investigating power series expansions in closed forms and integral forms of Lame function in the algebraic form and its asymptotic behaviors analytically.

VI. SERIES "SPECIAL FUNCTIONS AND THREE TERM RECURRENCE FORMULA (3TRF)"

This paper is 5th out of 10.

1. "Approximative solution of the spin free Hamiltonian involving only scalar potential for the $q-\bar{q}$ system"[23] - In order to solve the spin-free Hamiltonian with light quark masses we are led to develop a totally new kind of special function theory in mathematics that generalize all existing theories of confluent hypergeometric types. We call it the Grand Confluent Hypergeometric Function. Our new solution produces previously unknown extra hidden quantum numbers relevant for description of supersymmetry and for generating new mass formulas.

2. "Generalization of the three-term recurrence formula and its applications"[24] - Generalize three term recurrence formula in linear differential equation. Obtain the exact solution

of the three term recurrence for polynomials and infinite series.

3. "The analytic solution for the power series expansion of Heun function"[25] - Apply three term recurrence formula to the power series expansion in closed forms of Heun function (infinite series and polynomials) including all higher terms of A_ns.

4. "Asymptotic behavior of Heun function and its integral formalism",[26] - Apply three term recurrence formula, derive the integral formalism, and analyze the asymptotic behavior of Heun function (including all higher terms of A_ns).

5. "The power series expansion of Mathieu function and its integral formalism",[27] - Apply three term recurrence formula, analyze the power series expansion of Mathieu function and its integral forms.

6. "Lame equation in the algebraic form"[28] - Applying three term recurrence formula, analyze the power series expansion of Lame function in the algebraic form and its integral forms.

7. "Power series and integral forms of Lame equation in the Weierstrass's form and its asymptotic behaviors"[29] - Applying three term recurrence formula, derive the power series expansion of Lame function in the Weierstrass's form and its integral forms.

8. "The generating functions of Lame equation in the Weierstrass's form"[30] - Derive the generating functions of Lame function in the Weierstrass's form (including all higher terms of A_n's). Apply integral forms of Lame functions in the Weierstrass's form.

9. "Analytic solution for grand confluent hypergeometric function"[31] - Apply three term recurrence formula, and formulate the exact analytic solution of grand confluent hypergeometric function (including all higher terms of A_n's). Replacing μ and $\varepsilon\omega$ by 1 and $-q$, transforms the grand confluent hypergeometric function into Biconfluent Heun function.

10. "The integral formalism and the generating function of grand confluent hypergeometric function"[32] - Apply three term recurrence formula, and construct an integral formalism and a generating function of grand confluent hypergeometric function (including all higher terms of A_n's).

ACKNOWLEDGMENT

I thank Bogdan Nicolescu. The discussions I had with him on number theory was of great joy.

REFERENCES

[1] Choun, Y.S., “Generalization of the three-term recurrence formula and its applications,” arXiv:1303.0806

[2] Choun, Y.S. and Catto, S., “Approximative solution of the spin free Hamiltonian involving only scalar potential for the $q-\bar{q}$ system,” arXiv:1302.7309

[3] McLachlan, N. W., *Theory and application of Mathieu functions*, (Clarendon Press, Oxford, England, 1947).

[4] Gutirrez-Vega, J. C., Rodrguez-Dagnino,R. M., Meneses-Nava, M. A. and Chvez-Cerda, S., “Mathieu functions, a visual approach,” Amer. J. Phys. **71**(3), 233242(2003).

[5] Daymond, S. D., “The principal frequencies of vibrating systems with elliptic boundaries,” Quart. J. Mech. Appl. Math. **8**(3), 361372(1955).

[6] Troesch, B. A. and Troesch, H. R., “Eigenfrequencies of an elliptic membrane,” Math. Comp. **27**(124), 755765(1973).

[7] Alhargan, F. and Judah, S., “A general mode theory for the elliptic disk microstrip antenna,” IEEE Trans. Antennas and Propagation **43**(6), 560568(1995).

[8] Shen, L., “The elliptical microstrip antenna with circular polarization,” IEEE Trans. Antennas and Propagation **29**(1), 9094(1981).

[9] Bhattacharyya, A. and Shafai, L., “Theoretical and experimental investigation of the elliptical annual ring antenna,” IEEE Trans. Antennas and Propagation **36**(11), 15261530(1988).

[10] Ragheb, H. A., Shafai, L. and Hamid, M., “Plane wave scattering by a conducting elliptic cylinder coated by a nonconfocal dielectric,” IEEE Trans. Antennas and Propagation **39**(2), 218223(1991).

[11] Aliev, A.N., Hortacsu, M., Kalayci, J., and Nutku, Y., “Gravitational instantons from minimal surfaces,” Class. Quantum Grav. **16**, 631(1999).

[12] M. Hortacsu, “Heun Functions and their uses in Physics,” arXiv:1101.0471

[13] Frenkel, D. and Protugal, R., “Algebraic Methods to Compute Mathieu Functions,” J. Phys. A: Math. Gen. **34**, 3541(2001).

[14] Mathieu, E., “Mémoire sur Le Mouvement Vibratoire d’une Membrane de forme Elliptique,” J. Math. Pure Appl. **13**, 137 (1868).

[15] Sips, R., “Représentation asymptotique des fonctions de Mathieu et des fonctions d’onde sphéroidales,” Trans. Amer. Math. Soc. **66**, 93(1949).

[16] Whittaker, E.T.,“On the General Solution of Mathieu’s Equation,” Proc. Edinburgh Math. Soc. **32**, 75(1914).

[17] Wang, Z.W. and Guo, D.R., *Special functions*, (World Scientific Publishing Co Pte Ltd, New Jersey, 1989) p.610.

[18] Birkandan, T., Hortacsu, M.,“Examples of Heun and Mathieu functions as solutions of wave equations in curved spaces,”J. Phys. A: Math. Theor. **40**, 1105-1116(2007).

[19] Birkandan, T., Hortacsu, M.,“Dirac equation in the background of the Nutku helicoid metric,” J. Phys. A: Math. Theor. **48**, 092301(2007).

[20] Newman, E.T. and Penrose, R., “An Approach to Gravitational Radiation by a Method of Spin Coefficients,” J. Math. Phys. **3**, 566768(1962).

[21] Connor, J. N. L., Uzer, T., Marcus, R. A. and Smith, A. D., “Eigenvalues of the Schrdinger equation for a periodic potential with nonperiodic boundary conditions: A uniform semiclassical analysis,” J. Chem. Phys. **80**(10), 5095-6006(1984)

[22] Zlatev, I., Huey, G. and Steinhardt, P.J., “Parametric Resonance in an Expanding Universe,” Phys. Rev. D**57**, 2152-2157(1998).

[23] Link to arXiv:1302.7309.

[24] Link to arXiv:1303.0806.

[25] Link to arXiv:1303.0830.

[26] Link to arXiv:1303.0876.

[27] Link to arXiv:1303.0820.

[28] Link to arXiv:1303.0873.

[29] Link to arXiv:1303.0878.

[30] Link to arXiv:1303.0879.

[31] Link to arXiv:1303.0813.

[32] Link to arXiv:1303.0819.